


NAVAL POSTGRADUATE SCHOOL

Monterey, California



PRELIMINARY RESULTS FROM THE ANALYSIS OF
WIND COMPONENT ERROR
JULY DATA

Donald P. Gaver
//
Patricia A. Jacobs

November 1992

Approved for public release; distribution is unlimited.

Prepared for:
Naval Research Laboratory-West
Monterey, CA 93943-5006

Fed Docs
D 208.14/2 NPS-OR-93-003

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIFORNIA

Rear Admiral R. W. West, Jr.
Superintendent

Harrison Shull
Provost

This report was prepared in conjunction with research funded by the
Naval Research Laboratory-West, Monterey, California.

This report was prepared by:

UNCLASSIFIED
 SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
 OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NPSOR-93-003		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Naval Postgraduate School	6b. OFFICE SYMBOL (If applicable) OR	7a. NAME OF MONITORING ORGANIZATION Naval Research Laboratory-West	
6c. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		7b. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943-5006	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Research Laboratory-West	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Monterey, CA 93943-5006		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) <i>Preliminary Results from the Analysis of Wind Component Error-July Data</i>			
12. PERSONAL AUTHOR(S) Donald P. Gaver and Patricia A. Jacobs			
13a. TYPE OF REPORT Technical Report	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, month day) November 1992	15. PAGE COUNT 107
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Gaussian model with log-linear scale parameter; maximum likelihood; Newton's method	
FIELD	GROUP SUB-GROUP		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) <p>Estimation of mean square prediction error of wind components is required in the optimal interpolation (OI) process in numerical prediction of atmospheric variables. Statistical models with log-linear scale parameters which include covariates are described for the prediction error. Data from February April and July of 1991 are used to fit the model parameters and to study the predictive ability of the models. This preliminary investigation indicates that observational and first guess wind components can be helpful in predicting mean square prediction error for wind components. The predictions using observational winds appear to be better at the 850 mb level. The predictions using first guess winds appear to be better at the 250 mb level.</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL D. P. Gaver		22b. TELEPHONE (Include Area Code) (408) 646-2605	2c. OFFICE SYMBOL OR/Gv

PRELIMINARY RESULTS FROM THE ANALYSIS OF
WIND COMPONENT ERROR
JULY DATA

by

P. A. Jacobs and D. P. Gaver

TABLE OF CONTENTS

0. INTRODUCTION.....	1
1. THE MODELS.....	3
One Variable Models.....	3
2. ESTIMATION OF PARAMETERS.....	4
3. THE DATA ANALYSIS—JULY DATA.....	4
3.1 Observed Wind Covariate Models.....	4
3.2 First-guess Wind Covariate Models.....	12
4. A COMPARISON OF MODELS FOR THE MONTHS OF FEBRUARY, APRIL, AND JULY.....	18
4.1 Observed Wind Covariate Models.....	19
4.2 First-guess Wind Covariate Models.....	24
4.3 Conclusions.....	31
REFERENCES.....	56
APPENDIX A.....	57
APPENDIX B.....	70

PRELIMINARY RESULTS FROM THE ANALYSIS OF
WIND COMPONENT ERROR
JULY DATA

by

P. A. Jacobs and D. P. Gaver

0. INTRODUCTION

Numerical meteorological models are used to assist in the prediction of weather. Each run of a numerical model produces forecasts of meteorological variables which are used as preliminary predictions of the future values of these variables. These initial predictions are referred to as **first-guess** values. In this paper first-guess values will refer to the most recent 12 hour forecasts.

In certain areas of the world, observations of the values of forecasted variables become available. In our case the observations become available 12 hours after the first-guess values are computed. Prior to the next run of the numerical model a multivariate optimal interpolation analysis updates a first-guess value of a variable by adding to it a weighted observed value of the variable if it is available. The weight multiplying the observed value depends on estimates of the squared error of the first-guess value and the squared error of the observation; cf. Goerss et al. [1991, a, b]. Thus it is of importance to predict such first-guess squared errors.

The general problem of modeling and predicting mean square errors is important but not widely studied; see Davidian and Carroll (1987), Nelder and Lee (1992), Aitken (1987), and McCullagh and Nelder (1983). In the next

section statistical models for the error of the first-guess are introduced. The models assume the error of the first-guess has mean 0 but has a scale parameter that is log-linear with suitable covariates, i.e. explanatory or regression variables.

Results are reported concerning the estimation of model parameters, and model cross-validation and predictive ability for u, v wind component data from the month of July 1991. The data consist of measurements and 12 hour forecasts (first-guess values) at the 850 mb, 500 mb and 250 mb levels from 93 stations in North America, 25N–75N. The forecasts are produced using the NOGAPS Spectral Forecast Model; cf. Hogan et al. (1991). Each station has measurement and first-guess values for every 12 hours; there are some missing observations and suspicious values of wind components equal to 0. These missing and questionable values are deleted from the data set. The measurement values (if available) are subtracted from first-guess values to obtain observations of the error of the first-guess value. The results appear in Sections 3 and 4 and in Appendices A, B.

The results indicate that estimates of the variance of the error of first-guess wind components can be improved by using covariates which are functions of the wind components. Covariates using observed values of the wind components appear to have more predictive ability than those using first-guess values. Further exploratory work is needed to determine the degree with which these statistical results can be used to improve the forecasting ability of the numerical model.

1. THE MODELS

Fix a location. Let

$U_0(t)$ = observed u -wind component at time t

$U_f(t)$ = first-guess u -wind component at time t

$V_0(t)$ = observed v -wind component at time t

$V_f(t)$ = first-guess v -wind component at time t

$$r(t) = \left[(U_0(t) - U_0(t-1))^2 + (V_0(t) - V_0(t-1))^2 \right]^{\frac{1}{2}}$$

$$s(t) = \left[U_0(t)^2 + V_0(t)^2 \right]^{\frac{1}{2}}$$

$$Y(t) = U_0(t) - U_f(t) \quad \text{or} \quad Y(t) = V_0(t) - V_f(t)$$

The variable $Y(t)$ is the first guess error. The variable $r(t)$ is a measure of the observed change in the wind. The variable $s(t)$ is the observed wind speed.

The models considered are as follows:

One Variable Models

1. $\{Y(t)\}$ are independent normally distributed random variables with mean 0 and variance

$$\sigma_1^2(1;t) = \exp\{\alpha_1(1) + \beta_1(1)r(t)\}. \quad (1)$$

2. $\{Y(t)\}$ are independent normally distributed random variables with mean 0 and variance

$$\sigma_1^2(2;t) = \exp\{\alpha_1(2) + \beta_1(2)s(t)\}. \quad (2)$$

Two Variable Model

3. $\{Y(t)\}$ are independent normally distributed random variables with mean 0 and variance

$$\sigma_2^2(t) = \exp\{\alpha + \beta_1 r(t) + \beta_2 s(t)\}. \quad (3)$$

Independence Assumption

The first guess errors at different locations are independent. The parameters in the variance models do not depend on location.

2. ESTIMATION OF PARAMETERS

The model parameters are estimated by maximum likelihood. A system of equations is obtained by setting the first partial derivative with respect to each parameter of the log likelihood function equal to zero. The system of equations is solved numerically using Newton's method to obtain the maximum likelihood estimates. The procedure for the normal models above is given in Appendix A of Jacobs and Gaver [1991].

3. THE DATA ANALYSIS—JULY DATA

3.1 Observed Wind Covariate Models

In this subsection we report an assessment of the goodness of fit and cross-validation for the normal models (1)–(3) using observational wind components as covariates. There are six analyses; one for the u -wind component (respectively v -wind component) for each pressure level. Once missing values and suspicious wind values of 0 are deleted there are 3519 data values at the 850 mb level, 3833 values at the 500 mb level and 3830 values at the 250 mb level. Each analysis proceeds along the same lines. In what follows by data we mean triples $\{y(t), r(t), s(t)\}$.

In each analysis the data are randomly divided into two sets called DA and DB without regard to the values of the data.

The maximum likelihood parameter estimates for each model (1)–(3) are obtained for each set DA and DB and for all the data. The parameter estimates and their estimated standard errors (computed from the second partial derivatives of the likelihood evaluated at the estimates) appear in Table 1. Note that all the estimates are positive. Hence increased $r(t)$ and/or $s(t)$ values are associated with increased variance of the first guess value. This is plausible physically, since a large value of $r(t)$ is indicative of a change in the atmosphere and a large value of $s(t)$ is indicative of an active location in the atmosphere. The estimated variances $\sigma_1^2(1,t)$, $\sigma_1^2(2,t)$, $\sigma_2^2(t)$, are computed for the parameters estimated from DA and DB using (1)–(3) for each data point in DA and DB.

The models are for the variances of the observations rather than the observations themselves. One possible procedure to informally assess goodness-of-fit and cross-validate the models is by binning the data. To assess models (1) and (3) the data $(y(t), r(t), s(t))$ are binned into 10 bins based on ordering the values of $r(t)$ from smallest to largest. The data in the first bin correspond to the smallest values of $r(t)$; the data in the 10th bin correspond to the largest values of $r(t)$. Each bin contains about $\frac{1}{10}$ th of the data with the 10th bin containing a few more data. The averages of the estimated variances for models (1) and (3) are computed for each bin. The average $y(t)^2$ is also computed for each bin.

To assess models (2) and (3) the same procedure is used but the binning is based on the values of $s(t)$.

**TABLE 1. NORMAL MODELS
JULY DATA PARAMETER ESTIMATES
(STANDARD ERRORS)
OBSERVED WIND COVARIATES**

Pressure Level	Wind Comp.	Data Set	One-variate Models				Two-variate Models		
			$r(t)$		$s(t)$		$\log \text{MSE} = \alpha + \beta_1 r(t) + \beta_2 s(t)$		
			α	β	α	β	α	β_1	β_2
850	u	A	1.47 (0.06)	0.11 (0.008)	1.50 (0.06)	0.09 (0.007)	1.25 (0.06)	0.09 (0.01)	0.05 (0.009)
		B	1.42 (0.06)	0.12 (0.010)	1.41 (0.06)	0.10 (0.008)	1.13 (0.07)	0.08 (0.01)	0.07 (0.009)
		ALL	1.45 (0.04)	0.11 (0.006)	1.46 (0.04)	0.09 (0.005)	1.20 (0.05)	0.08 (0.007)	0.06 (0.006)
850	v	A	1.58 (0.06)	0.10 (0.008)	1.53 (0.06)	0.09 (0.008)	1.36 (0.07)	0.07 (0.01)	0.05 (0.009)
		B	1.51 (0.06)	0.11 (0.009)	1.52 (0.06)	0.10 (0.008)	1.21 (0.007)	0.09 (0.01)	0.06 (0.009)
		ALL	1.55 (0.04)	0.11 (0.006)	1.53 (0.04)	0.09 (0.006)	1.29 (0.05)	0.08 (0.007)	0.06 (0.006)
500	u	A	1.37 (0.06)	0.12 (0.008)	1.54 (0.06)	0.06 (0.005)	1.12 (0.07)	0.10 (0.008)	0.03 (0.005)
		B	1.45 (0.06)	0.10 (0.009)	1.66 (0.06)	0.04 (0.005)	1.24 (0.07)	0.09 (0.009)	0.02 (0.005)
		ALL	1.40 (0.04)	0.11 (0.006)	1.58 (0.04)	0.05 (0.005)	1.17 (0.05)	0.10 (0.006)	0.03 (0.004)
500	v	A	1.53 (0.06)	0.09 (0.009)	1.74 (0.06)	0.03 (0.005)	1.35 (0.08)	0.08 (0.009)	0.02 (0.005)
		B	1.45 (0.06)	0.11 (0.009)	1.59 (0.06)	0.05 (0.005)	1.20 (0.07)	0.09 (0.009)	0.03 (0.005)
		ALL	1.49 (0.04)	0.10 (0.006)	1.66 (0.05)	0.04 (0.004)	1.27 (0.05)	0.09 (0.007)	0.03 (0.004)
250	u	A	2.41 (0.06)	0.06 (0.005)	2.50 (0.06)	0.03 (0.003)	2.13 (0.07)	0.05 (0.005)	0.02 (0.003)
		B	2.42 (0.06)	0.06 (0.005)	2.53 (0.06)	0.03 (0.003)	2.13 (0.07)	0.05 (0.005)	0.02 (0.003)
		ALL	2.41 (0.04)	0.06 (0.003)	2.52 (0.04)	0.03 (0.002)	2.13 (0.05)	0.05 (0.004)	0.02 (0.002)
250	v	A	2.41 (0.06)	0.07 (0.005)	2.47 (0.06)	0.03 (0.003)	2.08 (0.07)	0.05 (0.005)	0.02 (0.003)
		B	2.49 (0.06)	0.05 (0.005)	2.39 (0.06)	0.03 (0.003)	2.16 (0.07)	0.04 (0.005)	0.02 (0.003)
		ALL	2.44 (0.04)	0.06 (0.003)	2.43 (0.04)	0.03 (0.002)	2.12 (0.05)	0.05 (0.003)	0.02 (0.002)

$$r(t) = [((u(t) - u(t-1))^2 + (v(t) - v(t-1))^2)]^{1/2}$$

$$s(t) = [u(t)^2 + v(t)^2]^{1/2}$$

NOTE: Data are divided into two sets randomly without regard to data values. One set is called A; the other is called B.

Figures 1-24 present graphs of the log [average $y(t)^2$] in each bin versus log [average estimated variance] in each bin for models (1) and (3) and models (2) and (3). Figures 1, 5, 9, 13, 17, 21 (respectively 2, 6, 10, 14, 18, 22) show the logarithm of the average of the $y(t)^2$ values of DA (respectively DB) versus the logarithm of the average of the estimated variances for each bin using the estimated parameters from DA (respectively DB). If a model were perfect, a point should be close to the 45° line shown.

Figures 3, 7, 11, 15, 19, 23, (respectively 4, 8, 12, 16, 20, 24) present graphs of log average $y(t)^2$ of DA (respectively DB) versus log average estimated variances using parameters estimated using data DB (respectively DA). Once again if the model were perfect, the points would be close to the 45° line.

Since the two-variate model (3) is shown with both one-variate models, it is possible to obtain some idea of the effect of the two different sets of bins on the log averages. In particular, the graphs corresponding to the 500 Mb height winds, Figures 9-16, show that the display of log averages can be quite sensitive to which variate is used to do the binning.

Keeping this binning sensitivity in mind, the figures suggest the following concerning the models using observed winds as covariates. It appears that of the two one-variate models, model (1) which uses $r(t)$ as the covariate is the better. The two-variate model (3) appears somewhat better than model (1). If wind speed is used as the single covariate, it appears to overstate the variance; the addition of the second covariate $r(t)$ in this case seems to tend to make the estimated variance smaller and bring the log [average predicted variance] in a bin closer to the log average y^2 in the bin.

Another way to assess goodness of fit and to cross validate is to evaluate the log-likelihood for the different models at the parameter estimates. Larger values of the log-likelihood suggest better model fit; cf. Cox and Hinkley [1974].

Table 2 presents the values of the log-likelihood at the parameter estimates up to addition and multiplication of constants for the parameter estimates of Table 1; the function being evaluated is

$$\tilde{\ell} = -n\alpha - \sum_{i=1}^n \underline{x}_i \underline{\beta} - \sum_{i=1}^n y_i^2 \exp\{-\alpha + \underline{x}_i \underline{\beta}\}. \quad (4)$$

where $\underline{x}_i \underline{\beta} = \sum_j x_{ij} \beta_j$. The values of $\tilde{\ell}$ are presented for data DA (respectively DB) using the $\hat{\beta}_j$ parameters fit using DA (respectively DB); these are values assessing goodness of fit; since maximum likelihood is the estimation procedure, the largest value of $\tilde{\ell}$ in each of these two rows is the one corresponding to the two-variate model. Values of $\tilde{\ell}$ are also presented for data DA (respectively DB) using the parameters fit using DB (respectively DA); these are values assessing cross-validation. The underlined value in each row is the maximum value in that row; the corresponding model provides the best model fit. The bold italicized value in each row is the maximum value for the two one-variate models; the corresponding one-variate model provides the best model fit between the two one-variate models.

The models considered in Table 2 are models (1)-(3) and the model that the $\{Y_i\}$ are independent normal with mean 0 and variance not a function of the covariates; that is,

TABLE 2. NORMAL MODELS
JULY DATA
OBSERVED WIND COVARIATES
LOG-LIKELIHOOD

Pressure Level	Wind Comp.	Data Set	Model	One-variate Models			Two-variate Models
				Constant	$r(t)$	$s(t)$	
850	u	A	A	-5727.0	-5424.4	-5468.8	<u>-5386.5</u>
		B	B	-5337.4	-5362.1	-5379.5	<u>-5306.5</u>
		B	A	-5546.9	-5363.6	-5381.8	<u>-5310.7</u>
		A	B	-5737.2	-5425.9	-5471.5	<u>-5391.2</u>
850	v	A	A	-5680.1	-5486.5	-5500.9	<u>-5448.5</u>
		B	B	-5693.0	-5504.0	-5540.3	<u>-5450.2</u>
		B	A	-5693.1	-5506.9	-5542.6	<u>-5457.8</u>
		A	B	-5680.1	-5489.7	-5503.4	<u>-5456.6</u>
500	u	A	A	-6237.3	-5909.8	-6049.7	<u>-5871.5</u>
		B	B	-5958.0	-5821.1	-5892.6	<u>-5795.8</u>
		B	A	-5977.0	-5827.4	-5912.7	<u>-5802.1</u>
		A	B	-6258.2	-5918.7	-6076.9	<u>-5879.9</u>
500	v	A	A	-6023.9	-5904.1	-5976.2	<u>-5889.2</u>
		B	B	-6193.6	-5997.8	-6072.8	<u>-5961.6</u>
		B	A	-6201.7	-6005.0	-6090.1	<u>-5971.2</u>
		A	B	-6031.5	-5910.2	-5990.6	<u>-5897.3</u>
250	u	A	A	-7893.9	-7680.0	-7762.9	<u>-7631.9</u>
		B	B	-7981.7	-7760.4	-7829.5	<u>-7703.2</u>
		B	A	-7983.7	-7760.7	-7830.2	<u>-7703.4</u>
		A	B	-7895.9	-7680.3	-7763.6	<u>-7632.1</u>
250	v	A	A	-8025.3	-7770.6	-7850.5	<u>-7710.6</u>
		B	B	-7758.0	-7611.7	-7622.3	<u>-7554.3</u>
		B	A	-7775.8	-7624.4	-7639.0	<u>-7569.5</u>
		A	B	-8044.9	-7786.3	-7868.7	<u>-7729.5</u>

$$\sigma_i^2(t) = e^\alpha \quad (\text{Constant variance}). \quad (5)$$

The two-variate model (3) maximizes the cross-validation values of $\tilde{\ell}$ for data DA (respectively DB) with a model using parameters fit using DB (respectively DA). This suggests that both $r(t)$ and $s(t)$ together have predictive ability.

For the one-variate models (1) and (2) the cross-validation values of $\tilde{\ell}$ for DA (respectively DB) using the parameters fit using DB (respectively DA) are maximized when $r(t)$ is the variable for all cases. This suggests that $r(t)$ by itself has better predictive value than the wind speed $s(t)$ by itself. The goodness of fit values of $\tilde{\ell}$ for the one-variate models using DA (respectively DB) have a higher value of $\tilde{\ell}$ associated with $r(t)$ the majority of the time. This suggests that $r(t)$ by itself provides a better description of the data than $s(t)$ by itself.

Comparing the value of $\tilde{\ell}$ for the model with constant variance (5), $\tilde{\ell}_c$, for DA (respectively DB) fit using DA (respectively DB) with the corresponding cross-validation value of $\tilde{\ell}$ for DA (respectively DB) using models (2), (3) fit using DB (respectively DA) indicates the following. The values of $\tilde{\ell}$ for models (1), (2) and (3) fit with the other half of the data are larger than the corresponding value $\tilde{\ell}_c$ for the constant variance model fit using the data to be modeled. This indicates that both models (2) and (3) fit with the other half of the data describe the data better than the best constant variance model (5) fit with the same data it is used to summarize.

Table 3 presents values of the fraction of increase in $\tilde{\ell}$, $(\tilde{\ell} - \tilde{\ell}_c) / |\tilde{\ell}_c|$, where $\tilde{\ell}_c$ is the maximum value of $\tilde{\ell}$ for the constant variance model (5) fit using data DA (respectively DB) compared to the cross-validation value of $\tilde{\ell}$ for DA

(respectively DB) using models (1)-(3) fit using DB (respectively DA). Large values of the fraction will indicate better model predictive ability. Note that the fraction of increase tends to become larger for higher pressure levels. This behavior suggests that if winds from one pressure level are to be used to estimate the variance of the first guess, it should be the 850 mb level. Comparison of the values for the two one-variate models once again suggests

TABLE 3. JULY OBSERVED WIND COVARIATES
FRACTION OF INCREASE $(\bar{\ell} - \bar{\ell}_c) / |\bar{\ell}_c|$

Pressure Level	Wind Comp.	Data Set	Model	One-variate Models		Two-variate Models
				$r(t)$	$s(t)$	
850	u	B	A	0.03	0.03	0.04
		A	B	0.05	0.04	0.06
	v	B	A	0.04	0.03	0.04
		A	B	0.03	0.03	0.04
500	u	B	A	0.02	0.008	0.03
		A	B	0.05	0.03	0.06
	v	B	A	0.03	0.02	0.04
		A	B	0.02	0.01	0.02
250	u	B	A	0.03	0.02	0.03
		A	B	0.03	0.02	0.03
	v	B	A	0.02	0.02	0.02
		A	B	0.03	0.02	0.04

that the one-variate model using $r(t)$ has the greater predictive ability. Once again the two-variate model appears to have the most predictive ability.

To further explore the predictive ability of the models using observed covariates, bootstrap experiments were conducted. Six bootstrap experiments were conducted; one for each u and v wind component at each pressure level.

Each experiment consists of 250 replications. For each replication the data are randomly divided into two sets independent of their values which we will call A and B. Models (1)-(3) are fit to each data set. The value of $\tilde{\ell}_c$, the value of $\tilde{\ell}$ for the constant variance model fit using the same data it is to describe, is computed for each data set. The value of $\tilde{\ell}$ for each data set is computed for each model (1)-(3) with parameters estimated using the other half of the data. The fraction of increase in $\tilde{\ell}$, $(\tilde{\ell}-\tilde{\ell}_c)/|\tilde{\ell}_c|$ is computed for each half of the data. Figures 1A-6A in Appendix A display histograms of $(\tilde{\ell}-\tilde{\ell}_c)/|\tilde{\ell}_c|$ for models using observed wind covariates. Each histogram includes the fractions for both A and B data sets. The histogram indicated that the models for the 850 mb level have the most predictive ability. Model (3) using both covariates appears to have somewhat better predictive ability. Of the two one-variate models model (1) using $r(t)$ as the covariate clearly has the better predictive ability.

3.2 First-guess Wind Covariate Models

In this section we report the results of using models (1)-(3) and (5) with first-guess winds as covariates; the two covariates considered are

$$r_f(t) = \left[\left(U_f(t) - U_f(t-1) \right)^2 + \left(V_f(t) - V_f(t-1) \right)^2 \right]^{\frac{1}{2}}$$

and

$$s_f(t) = \left[U_f(t)^2 + V_f(t)^2 \right]^{\frac{1}{2}}.$$

The first guess resultant wind $r_f(t)$ is a measure of forecasted change in the winds. The first guess wind speed $s_f(t)$ is a measure of forecasted activity

in the atmosphere. Since observed winds are not available over a great portion of the earth, it is important to have models for predicting the variance of the first-guess values which involve the first-guess values which are always available.

Once missing values and suspicious 0 wind values are deleted, there are 3710 observations at the 850 mb level, 4208 observations at the 500 mb level, and 4132 observations at the 250 mb level. The analysis is the same as in the previous subsection. The data sets DA and DB are the same as those in the previous subsection in each case. The values of the parameter estimates with estimated standard errors appear in Table 4. Note that the estimates are all positive. Hence increased $r_f(t)$ and/or $s_f(t)$ is associated with higher variance of the first guess error. The corresponding values of $\tilde{\ell}$ appear in Table 5. Once again the underlined value of $\tilde{\ell}$ is the largest value in each row; the bold italicized value $\tilde{\ell}$ is the largest value between the two one-variate models.

In all cases the values of $\tilde{\ell}$ for the observed wind covariates are larger than those for the first-guess wind covariates. This suggests that the observed wind components have better predictive and descriptive value than the first-guess wind components.

**TABLE 4. NORMAL MODELS
PARAMETER ESTIMATES
(STANDARD ERROR)
FIRST-GUESS WIND COVARIATES**

Pressure Level	Wind Comp.	Data Set	One-variate Models				Two Variate Models		
			$r_f(t)$		$s_f(t)$		$\log \text{MSE} = \alpha + \beta_1 r_f(t) + \beta_2 s_f(t)$		
			α	β	α	β	α	β_1	β_2
850	<i>u</i>	A	2.04 (0.06)	0.05 (0.01)	2.13 (0.06)	0.02 (0.01)	2.01 (0.07)	0.05 (0.01)	0.005 (0.01)
		B	1.96 (0.06)	0.05 (0.01)	2.00 (0.06)	0.02 (0.009)	1.90 (0.07)	0.04 (0.01)	0.01 (0.009)
		ALL	2.00 (0.04)	0.05 (0.009)	2.07 (0.04)	0.02 (0.006)	1.96 (0.05)	0.05 (0.009)	0.009 (0.007)
850	<i>v</i>	A	2.03 (0.06)	0.04 (0.01)	1.96 (0.06)	0.03 (0.009)	1.91 (0.07)	0.02 (0.01)	0.03 (0.009)
		B	1.94 (0.06)	0.07 (0.01)	1.93 (0.06)	0.05 (0.008)	1.82 (0.07)	0.06 (0.01)	0.03 (0.009)
		ALL	1.98 (0.04)	0.06 (0.008)	1.94 (0.04)	0.04 (0.006)	1.86 (0.05)	0.04 (0.009)	0.03 (0.007)
500	<i>u</i>	A	1.87 (0.05)	0.06 (0.01)	1.75 (0.06)	0.03 (0.005)	1.63 (0.07)	0.04 (0.01)	0.03 (0.005)
		B	1.99 (0.06)	0.04 (0.01)	1.89 (0.06)	0.03 (0.005)	1.82 (0.07)	0.02 (0.01)	0.02 (0.005)
		ALL	1.93 (0.04)	0.05 (0.007)	1.82 (0.04)	0.03 (0.003)	1.72 (0.05)	0.03 (0.008)	0.03 (0.004)
500	<i>v</i>	A	1.95 (0.05)	0.05 (0.01)	1.84 (0.06)	0.03 (0.005)	1.75 (0.07)	0.03 (0.01)	0.02 (0.005)
		B	1.99 (0.05)	0.04 (0.01)	2.01 (0.06)	0.01 (0.005)	1.91 (0.07)	0.04 (0.01)	0.009 (0.005)
		ALL	1.97 (0.04)	0.04 (0.007)	1.92 (0.04)	0.02 (0.004)	1.83 (0.05)	0.03 (0.007)	0.02 (0.004)
250	<i>u</i>	A	2.79 (0.06)	0.04 (0.006)	2.64 (0.06)	0.02 (0.003)	2.54 (0.06)	0.02 (0.007)	0.02 (0.003)
		B	3.00 (0.05)	0.03 (0.006)	2.92 (0.06)	0.02 (0.003)	2.84 (0.06)	0.02 (0.005)	0.01 (0.003)
		ALL	3.00 (0.04)	0.03 (0.004)	2.79 (0.04)	0.02 (0.002)	2.70 (0.05)	0.02 (0.006)	0.02 (0.002)
250	<i>v</i>	A	2.81 (0.05)	0.04 (0.006)	2.71 (0.06)	0.02 (0.003)	2.58 (0.07)	0.03 (0.006)	0.02 (0.003)
		B	2.77 (0.05)	0.05 (0.006)	2.70 (0.06)	0.02 (0.003)	2.51 (0.07)	0.04 (0.006)	0.02 (0.003)
		ALL	2.79 (0.04)	0.04 (0.004)	2.71 (0.04)	0.02 (0.002)	2.55 (0.05)	0.03 (0.004)	0.02 (0.002)

TABLE 5.
JULY LOG-LIKELIHOOD
FIRST-GUESS COVARIATES

Pressure Level	Wind Comp.	Data Set	Model	Constant	One-variate Models		Two-variate Models
					$r_f(t)$	$s_f(t)$	
850	u	A	A	-6018.5	-6000.0	-6014.4	<u>-5999.7</u>
		B	B	-5857.1	-5840.3	-5848.5	<u>-5837.9</u>
		B	A	-5863.9	-5847.4	-5856.9	<u>-5846.3</u>
		A	B	-6025.7	<u>-6007.5</u>	-6023.6	-6009.0
850	v	A	A	-5900.1	-5890.2	-5884.7	<u>-5881.0</u>
		B	B	-6023.1	-5977.5	-5987.7	<u>-5967.3</u>
		B	A	-6027.2	-5990.4	-5994.1	<u>-5978.4</u>
		A	B	-5904.1	-5900.2	-5900.5	<u>-5889.8</u>
500	u	A	A	-6624.7	-6584.1	-6567.8	<u>-6550.7</u>
		B	B	-6683.1	-6669.1	-6653.3	<u>-6649.0</u>
		B	A	-6683.9	-6674.1	-6658.8	<u>-6656.9</u>
		A	B	-6625.5	-6589.5	-6573.2	<u>-6559.0</u>
500	v	A	A	-6658.3	-6636.2	-6622.8	<u>-6613.9</u>
		B	B	-6656.4	-6638.1	-6648.1	<u>-6635.4</u>
		B	A	-6656.4	<u>-6638.6</u>	-6655.4	-6643.6
		A	B	-6658.3	-6636.7	-6631.0	<u>-6623.1</u>
250	u	A	A	-8484.0	-8437.4	-8407.3	<u>-8392.6</u>
		B	B	-8693.2	-8670.7	-8660.6	<u>-8652.7</u>
		B	A	-8704.2	-8689.5	-8680.6	-8682.5
		A	B	-8494.3	-8454.1	-8431.0	<u>-8418.0</u>
250	v	A	A	-8453.7	-8411.9	-8399.7	<u>-8380.1</u>
		B	B	-8550.1	-8486.4	-8488.8	<u>-8451.2</u>
		B	A	-8552.4	-8491.0	-8490.1	<u>-8455.4</u>
		A	B	-8455.9	-8416.5	-8400.9	<u>-8384.2</u>

Table 5 also indicates the following results concerning models using first-guess wind covariates. Between the two one-variate models (1) and (2) the one-variate model using first-guess wind speed has the greater $\tilde{\ell}$ -value the majority of the time. This suggests that first-guess wind speed alone has somewhat better predictive and descriptive value than $r_f(t)$ alone. The cross-validation values of $\tilde{\ell}$ for data DA (respectively DB) using parameters fit with DB (respectively DA) are maximized in all cases except two for the two-variate model. This suggests that the two-variate model has better predictive ability.

Comparing the values of $\tilde{\ell}$, $\tilde{\ell}_c$, for DA (respectively DB) using the constant variance model (5) fit using DA (respectively DB) with the cross-validation value of $\tilde{\ell}$ for DA (respectively DB) using models (2), (3) fit using DB (respectively DA) indicates the following. The values of $\tilde{\ell}$ for models (1), (2) and (3) fit with the other half of the data are larger in all but two cases than the corresponding value $\tilde{\ell}_c$ for the constant variance model fit using the data to be modeled. This suggests that models (1)–(3) fit with the other half of the data describe the data somewhat better than the best constant variance model (5) fit with the data to be described.

Table 6 presents the fraction of increase in log-likelihood obtained by using models (1)–(3) fit using data DA (respectively DB) to describe data DB (respectively DA) compared to the value of the likelihood obtained by fitting the constant variance model (5) using data DB (respectively DA); Table 6 shows values of $\frac{\tilde{\ell} - \tilde{\ell}_c}{|\tilde{\ell}_c|}$. The results suggest that models using first guess wind have better predictive ability for lower pressure levels. Hence, if only one pressure level is to be used it is suggested that models for either the 500 mb level or 250 mb level be considered.

TABLE 6
 JULY DATA
 FRACTION OF INCREASE IN LOG-LIKELIHOOD
 FIRST GUESS COVARIATES

$$\frac{\bar{\ell} - \bar{\ell}_c}{|\bar{\ell}_c|}$$

Pressure Level	Wind Comp.	Data Set	Model	One-variate Models		Two-variate Models
				$r(t)$	$s(t)$	
850	u	B	A	0.002	0	0.002
		A	B	0.002	*	0.002
	v	B	A	0.005	0.005	0.007
		A	B	*	0.002	0.002
500	u	B	A	0.001	0.004	0.004
		A	B	0.005	0.008	0.010
	v	B	A	0.003	0.000	0.002
		A	B	0.003	0.004	0.005
250	u	B	A	0.000	0.000	0.001
		A	B	0.004	0.006	0.008
	v	B	A	0.007	0.007	0.011
		A	B	0.004	0.006	0.008

* $\bar{\ell}_c > \bar{\ell}$.

To further explore the predictive ability of the models using first guess covariates, bootstrap experiments were conducted. Six experiments were conducted, one for each u and v wind component at each pressure level. Each experiment consists of 250 replications. For each replication the data are randomly divided into two sets, independent of their values, which we will call A and B. Models (1)-(3) are fit to each data set. The value of $\bar{\ell}_c$, the value of $\bar{\ell}$ for the constant variance model fit using the same data it is to describe, is computed for each data set. The value of $\bar{\ell}$ for each data set is computed for each model (1)-(3) fit using the other half of the data. The fraction of increase in $\bar{\ell}$, $(\bar{\ell} - \bar{\ell}_c) / |\bar{\ell}_c|$, is computed for each half of the data. Figures 7A-12A in

Appendix A display histograms of $(\bar{\ell} - \bar{\ell}_c) / |\bar{\ell}_c|$ for models using first guess wind covariates. Each histogram includes the fractions for both A and B data sets. The histograms indicate that the models using first guess wind covariates do not have as much predictive ability as those using observed wind covariates. The first guess wind covariate models appear to have the most predictive ability at the 500 and 250 mb levels with the models at the 250 mb level being somewhat better. Model (3) using both first guess covariates appears to have the best predictive ability. Of the two one-variate models, Model (1) using $r_f(t)$ as the covariate has the better predictive ability. The predictive ability of the one-variate model using $s_f(t)$ is the most variable.

In summary, based on values of $\bar{\ell}$, when first-guess winds are used as covariates it appears that the two-variate model using first-guess wind speed at the 250 mb level is an attractive choice for predictive purposes. When observational winds are used as covariates, the two-variate model at the 850 mb level appears to have the best predictive value.

4. A COMPARISON OF MODELS FOR THE MONTHS OF FEBRUARY, APRIL, AND JULY

Results of a statistical analysis of the first-guess error field for the months of February 1991 and April 1991 are presented in Jacobs and Gaver (1991).

In this section we report results concerning the use of models fit with July data (respectively February or April) to predict February or April (respectively July) mean square first-guess error. These results give an indication of the possibility of using a model fit with one month's data to predict another month's data.

4.1 Observed Wind Covariate Models

In this subsection we report results for normal models (1)–(3) using observed wind components as covariates. There are six analyses; one for the u -wind component (respectively v -wind component) for each pressure level.

Table 7 shows the values of the parameter estimates and estimated standard errors for the February, April, and July data. The minor discrepancies with the values in Jacobs and Gaver (1991) are due to the deletion of the suspect 0 wind values from the data sets in this analysis. Table 8 shows the values of $\tilde{\ell}$ for February data (respectively July data) using parameters fit using February data (respectively July data). Values of $\tilde{\ell}$ are also presented for February (respectively July) data using parameters fit using July (respectively February) data. Once again, larger values of $\tilde{\ell}$ indicate better model fit. The underlined value in each row is the maximum value in that row. The bold italicized value in each row is the maximum value of $\tilde{\ell}$ for the two one-variate models.

The values of $\tilde{\ell}$ for February data (respectively July data) using parameters fit using July data (respectively February data) are maximized by the two-variate model in all but one case; between the two one-variate models $\tilde{\ell}$ is the maximized for the model involving $s(t)$ except in 3 cases.

Comparing the value of $\tilde{\ell}$, $\tilde{\ell}_c$, for the model of constant variance (5) for February (respectively July) data using parameters estimated from February (respectively July) data with that for the prediction value of $\tilde{\ell}$ for the models (2)–(3) for February (respectively July) data using parameters estimated from July (respectively February) data indicate the following. The values of $\tilde{\ell}$ for

**TABLE 7. PARAMETER ESTIMATES
(STANDARD ERRORS)
OBSERVED WIND COVARIATES**

Pressure Level	Wind Comp.	Data Set	One-Variate Models				Two-variate Models		
			$r(t)$		$s(t)$		$\log \text{MSE} = \alpha + \beta_1 r(t) + \beta_2 s(t)$		
			α	β	α	β	α	β_1	β_2
850	u	July	1.45 (0.04)	0.11 (0.006)	1.46 (0.04)	0.09 (0.005)	1.20 (0.05)	0.08 (0.007)	0.06 (0.006)
		Apr.	1.86 (0.04)	0.09 (0.005)	1.68 (0.04)	0.009 (0.004)	1.42 (0.05)	0.05 (0.005)	0.07 (0.005)
		Feb.	2.09 (0.04)	0.05 (0.005)	1.92 (0.05)	0.05 (0.004)	1.74 (0.05)	0.03 (0.005)	0.04 (0.004)
	v	July	1.55 (0.04)	0.11 (0.006)	1.53 (0.04)	0.09 (0.006)	1.29 (0.05)	0.08 (0.007)	0.06 (0.006)
		Apr.	1.84 (0.04)	0.09 (0.005)	1.68 (0.04)	0.09 (0.005)	1.43 (0.05)	0.06 (0.005)	0.07 (0.005)
		Feb.	2.15 (0.04)	0.05 (0.004)	1.71 (0.05)	0.07 (0.004)	1.50 (0.05)	0.03 (0.005)	0.06 (0.004)
500	u	July	1.40 (0.04)	0.11 (0.006)	1.58 (0.04)	0.05 (0.003)	1.17 (0.05)	0.10 (0.006)	0.03 (0.004)
		Apr.	2.12 (0.04)	0.06 (0.003)	2.22 (0.04)	0.03 (0.002)	1.81 (0.05)	0.05 (0.003)	0.02 (0.002)
		Feb.	2.22 (0.04)	0.05 (0.003)	2.40 (0.05)	0.02 (0.002)	2.02 (0.05)	0.05 (0.003)	0.01 (0.002)
	v	July	1.49 (0.04)	0.10 (0.006)	1.66 (0.05)	0.04 (0.004)	1.27 (0.05)	0.09 (0.007)	0.03 (0.004)
		Apr.	2.03 (0.04)	0.06 (0.003)	1.99 (0.04)	0.04 (0.002)	1.68 (0.05)	0.05 (0.004)	0.02 (0.003)
		Feb.	2.28 (0.04)	0.04 (0.003)	2.32 (0.05)	0.02 (0.002)	2.02 (0.05)	0.04 (0.003)	0.01 (0.002)
250	u	July	2.42 (0.04)	0.06 (0.003)	2.52 (0.04)	0.03 (0.002)	2.13 (0.05)	0.05 (0.004)	0.02 (0.002)
		Apr.	2.76 (0.04)	0.04 (0.002)	2.67 (0.04)	0.03 (0.001)	2.30 (0.05)	0.04 (0.002)	0.02 (0.002)
		Feb.	3.02 (0.04)	0.04 (0.002)	2.56 (0.04)	0.03 (0.001)	2.28 (0.06)	0.03 (0.002)	0.03 (0.001)
	v	July	2.44 (0.04)	0.06 (0.003)	2.43 (0.04)	0.03 (0.002)	2.12 (0.05)	0.05 (0.008)	0.02 (0.002)
		Apr.	2.75 (0.04)	0.04 (0.002)	2.64 (0.04)	0.03 (0.001)	2.28 (0.05)	0.03 (0.002)	0.02 (0.002)
		Feb.	3.00 (0.04)	0.003 (0.002)	2.43 (0.05)	0.03 (0.001)	2.23 (0.06)	0.02 (0.002)	0.03 (0.001)

$$r(t) = [((u(t) - u(t-1))^2 + (v(t) - v(t-1))^2)]^{1/2}$$

$$s(t) = [u(t)^2 + v(t)^2]^{1/2}$$

TABLE 8. NORMAL MODELS
VALUES OF LOG-LIKELIHOOD
OBSERVED WIND COVARIATES
FEBRUARY AND JULY

Pressure Level	Wind Comp.	Data Set	Model	Constant	One-Variate Models		Two-variate Models
					$r(t)$	$s(t)$	
850	u	July	July	-11269.3	-10787.3	-10849.5	-10695.2
		Feb.	Feb.	-13211.6	-13071.2	-13023.0	-12964.0
		Feb.	July	-13405.8	-13325.2	-13138.1	<u>-13116.5</u>
		July	Feb.	-11417.3	-11017.6	-10963.0	<u>-10826.0</u>
	v	July	July	-11373.1	-10992.0	-11042.4	-10902.7
		Feb.	Feb.	-13333.7	-13204.4	-12992.0	-12957.9
		Feb.	July	-13531.8	-13446.7	-13018.8	-13078.6
		July	Feb.	-11523.8	-11200.2	-11059.6	<u>-10972.4</u>
500	u	July	July	-12205.2	-11734.6	-11953.8	-11670.9
		Feb.	Feb.	-16273.0	-15924.6	-16151.7	-15892.9
		Feb.	July	-17497.3	-16399.1	-16512.5	<u>-16216.4</u>
		July	Feb.	-12913.9	-12174.5	-12419.9	<u>-12014.1</u>
	v	July	July	-12221.4	-11905.2	-12056.9	-11855.2
		Feb.	Feb.	-15966.1	-15750.5	-15859.9	-15707.1
		Feb.	July	-16900.4	-16168.5	-16066.0	<u>-15997.7</u>
		July	Feb.	-12790.9	-12281.1	-12361.7	<u>-12103.1</u>
250	u	July	July	-15876.6	-15440.6	-15592.7	-15335.2
		Feb.	Feb.	-18771.3	-17773.0	-17619.9	-17413.1
		Feb.	July	-20206.9	-18045.1	-17657.2	<u>-17530.2</u>
		July	Feb.	-16742.6	-15713.3	-15609.6	<u>-15386.4</u>
	v	July	July	-15792.6	-15389.4	-15481.5	-15273.3
		Feb.	Feb.	-18095.0	-17366.6	-17227.4	-17062.1
		Feb.	July	-18953.0	-17603.0	-17227.8	<u>-17186.8</u>
		July	Feb.	-16366.7	-15608.7	-15481.6	<u>-15323.9</u>

models (2) and (3) fit with data from the other month are larger in the majority of the cases than the corresponding values of $\tilde{\ell}_c$ fit with the data of the same month. This suggests that models (2) and (3) fit using data from the other month have some predictive value over a model of constant variance fit using the data that is to be modeled.

Table 9 shows values of $\tilde{\ell}$ for April data (respectively July data) using parameters fit using April data (respectively July data). Values of $\tilde{\ell}$ are also presented for April data (respectively July data) using parameters fit using July data (respectively February data). The underlined value in each row is the maximum value in that row. The bold italicized value in each row is the maximum value of $\tilde{\ell}$ for the two one-variate models.

The values of $\tilde{\ell}$ for April data (respectively July data) using parameters fit using July data (respectively April data) are maximized by the two-variate model in all cases; between the two one-variate models $\tilde{\ell}$ is maximized in all but five cases for the model involving $r(t)$.

Comparing the value of $\tilde{\ell}$, $\tilde{\ell}_c$ for the model of constant variance (5) for April (respectively July) data using parameters estimated from April (respectively July) data with that for the prediction value of $\tilde{\ell}$ for the models (2)-(3) for April (respectively July) data using parameters estimated from July (respectively April) data indicate the following. The values of $\tilde{\ell}$ for models (2) and (3) fit with data from the other month are larger in the majority of the cases than the corresponding values of $\tilde{\ell}_c$ fit with the data of the same month. This suggests that models (2) and (3) fit using data from the other month have some predictive value over a model of constant variance fit using the data that is to be modeled.

TABLE 9. OBSERVED WIND COVARIATES
VALUE OF LOG-LIKELIHOOD
APRIL AND JULY

Pressure Level	Wind Comp.	Data Set	Model	Constant	One-Variate Models		Two-variate Models
					$r(t)$	$s(t)$	
850	u	July	July	-11269.3	-10787.3	-10849.5	-10695.2
		April	April	-13814.3	-13460.9	-13313.6	-13205.7
		April	July	-14081.0	-13592.6	-13369.6	<u>-13265.6</u>
		July	April	-11460.5	-10911.4	-10901.0	<u>-10743.5</u>
	v	July	July	-11373.1	-10992.0	-11042.4	-10902.7
		April	April	-13837.2	-13421.2	-13389.5	-13229.7
		April	July	-14067.2	-13490.0	-13423.9	<u>-13251.4</u>
		July	April	-11540.6	-11058.4	-11073.6	<u>-10920.7</u>
500	u	July	July	-12205.2	-11734.6	-11953.8	-11670.9
		April	April	-16262.1	-15875.3	-16055.1	-15775.5
		April	July	-17101.2	-16259.4	-16391.3	<u>-16020.9</u>
		July	April	-12714.1	-12074.3	-12272.1	<u>-11893.0</u>
	v	July	July	-12221.4	-11905.2	-12056.9	-11855.2
		April	April	-16476.6	-15698.2	-15843.3	-15584.2
		April	July	-17472.3	-15913.3	-16008.0	<u>-15703.2</u>
		July	April	-12807.2	-12095.5	-12198.1	<u>-11946.8</u>
250	u	July	July	-15876.6	-15440.6	-15592.7	-15335.2
		April	April	-20104.9	-17863.0	-18119.6	-17705.0
		April	July	-21601.8	-17954.3	-18144.5	<u>-17750.3</u>
		July	April	-16723.4	-15514.5	-15619.0	<u>-15357.4</u>
	v	July	July	-15792.6	-15389.4	-15481.5	-15273.3
		April	April	-18674.8	-17610.7	-17853.7	-17473.4
		April	July	-19096.9	-17691.2	-17884.1	<u>-17525.5</u>
		July	April	-16089.6	-15448.2	-15507.4	<u>-15296.4</u>

Table 10 shows the fraction of increase in $\tilde{\ell}$ of using a model with parameters estimated using another month to predict variance in the current month compared to using the best constant variance model fit with the current month. The results suggest that the models for other months do have some predictive ability. Models fit using April data appear to have more predictive ability for July than those fit using February data. The predictive ability appears greater at the 250 mb level.

4.2 First-guess Wind Covariate Models

In this section we report results for normal models (1)–(3) using first-guess wind components as covariates.

Table 11 shows the values of the parameter estimates and standard errors for February data, April data and July data. The minor discrepancies with values reported in Jacobs and Gaver (1991) are due to the deletion of suspicious 0 wind values from the data sets. Table 12 shows the values of $\tilde{\ell}$ for February data (respectively April data) using parameters estimated from February data (respectively July data). Values of $\tilde{\ell}$ are also presented for February data (respectively July data) using parameters estimated from July data (respectively February data). The underlined value in each row is the maximum value in that row. The bold italicized value in each row is the maximum value of $\tilde{\ell}$ for the two one-variate models.

The values of $\tilde{\ell}$ for the observed wind covariates are larger than those for the first-guess wind covariates in all cases. This suggests that the observed wind covariates provide better models of the data both in terms of goodness-of-fit and prediction.

TABLE 10. FRACTION OF INCREASE
IN LOG-LIKELIHOOD
 $(\bar{\ell} - \bar{\ell}_c) / |\bar{\ell}_c|$

Pressure Level	Wind Comp.	Data Set	Model	One-variate Models		Two-variate
				$r(t)$	$s(t)$	Models
850	u	July	Feb.	0.02	0.03	0.04
		July	Apr.	0.03	0.03	0.05
		Feb.	July	*	0.006	0.007
		Apr.	July	0.02	0.03	0.04
	v	July	Feb.	0.02	0.03	0.04
		July	Apr.	0.03	0.03	0.04
		Feb.	July	*	0.006	0.007
		Apr.	July	0.03	0.03	0.04
500	u	July	Feb.	0.003	*	0.02
		July	Apr.	0.01	*	0.03
		Feb.	July	*	*	*
		Apr.	July	0.00	*	0.01
	v	July	Feb.	*	*	0.01
		July	Apr.	0.01	0.002	0.02
		Feb.	July	*	*	*
		Apr.	July	0.03	0.03	0.05
250	u	July	Feb.	0.01	0.02	0.03
		July	Apr.	0.02	0.02	0.03
		Feb.	July	0.04	0.06	0.07
		Apr.	July	0.11	0.10	0.12
	v	July	Feb.	0.01	0.02	0.03
		July	Apr.	0.02	0.02	0.03
		Feb.	July	0.03	0.05	0.05
		Apr.	July	0.05	0.04	0.06

*: $\bar{\ell}_c$ (data described by model with constant variance estimated using same data)

> $\bar{\ell}$ (data described by model fit using other month)

**TABLE 11. FIRST GUESS WIND COVARIATES
PARAMETER ESTIMATES
(STANDARD ERRORS)**

Pressure Level	Wind Comp.	Data Set	One-variate Models				Two-variate Models		
			$r(t)$		$s(t)$		$\log \text{MSE} = \alpha + \beta_1 r(t) + \beta_2 s(t)$		
			α	β	α	β	α	β_1	β_2
850	u	July	2.00 (0.04)	0.05 (0.009)	2.07 (0.04)	0.02 (0.006)	1.96 (0.05)	0.045 (0.009)	0.009 (0.007)
		Apr	2.37 (0.04)	0.03 (0.006)	2.12 (0.04)	0.05 (0.004)	2.11 (0.05)	0.003 (0.007)	0.045 (0.004)
		Feb	2.47 (0.04)	0.01 (0.005)	2.25 (0.04)	0.03 (0.004)	2.27 (0.05)	-0.01 (0.005)	-0.03 (0.004)
850	v	July	1.98 (0.04)	0.06 (0.008)	1.94 (0.04)	0.04 (0.006)	1.86 (0.05)	0.04 (0.009)	0.03 (0.007)
		Apr	2.46 (0.04)	0.02 (0.006)	2.21 (0.04)	0.04 (0.004)	2.22 (0.05)	-0.002 (0.007)	0.04 (0.004)
		Feb	2.45 (0.04)	0.01 (0.005)	2.35 (0.04)	0.02 (0.003)	2.34 (0.04)	0.003 (0.005)	0.02 (0.004)
500	u	July	1.93 (0.04)	0.05 (0.007)	1.82 (0.04)	0.03 (0.003)	1.72 (0.05)	0.03 (0.008)	0.03 (0.004)
		Apr	2.51 (0.04)	0.03 (0.005)	2.25 (0.04)	0.03 (0.002)	2.14 (0.05)	0.02 (0.005)	0.03 (0.002)
		Feb	2.61 (0.04)	0.03 (0.004)	2.54 (0.05)	0.02 (0.002)	2.38 (0.05)	0.02 (0.004)	0.01 (0.002)
500	v	July	1.97 (0.04)	0.04 (0.007)	1.92 (0.04)	0.02 (0.004)	1.83 (0.05)	0.03 (0.007)	0.02 (0.004)
		Apr	2.33 (0.04)	0.06 (0.005)	1.96 (0.04)	0.05 (0.002)	1.76 (0.05)	0.03 (0.005)	0.04 (0.002)
		Feb	2.71 (0.04)	0.01 (0.004)	2.47 (0.05)	0.02 (0.002)	2.44 (0.05)	0.004 (0.004)	0.01 (0.002)
250	u	July	2.90 (0.04)	0.03 (0.004)	2.79 (0.04)	0.02 (0.002)	2.70 (0.05)	0.02 (0.005)	0.02 (0.002)
		Apr	4.01 (0.04)	-0.01 (0.004)	3.48 (0.05)	0.01 (0.002)	3.63 (0.06)	-0.02 (0.004)	0.02 (0.002)
		Feb	3.67 (0.04)	0.02 (0.003)	2.94 (0.05)	0.03 (0.002)	2.75 (0.06)	0.02 (0.03)	0.03 (0.001)
250	v	July	2.79 (0.04)	0.04 (0.004)	2.71 (0.04)	0.02 (0.002)	2.55 (0.05)	0.03 (0.004)	0.02 (0.002)
		Apr	3.27 (0.04)	0.03 (0.003)	2.80 (0.05)	0.03 (0.002)	2.68 (0.05)	0.01 (0.003)	0.02 (0.002)
		Feb	3.48 (0.04)	0.02 (0.003)	3.08 (0.05)	0.02 (0.002)	2.84 (0.06)	0.02 (0.003)	0.02 (0.001)

TABLE 12. FIRST-GUESS WIND COVARIATES
VALUE OF LOG-LIKELIHOOD
JULY AND FEBRUARY

Pressure Level	Wind Comp.	Data Set	Model	One-variate Models			Two-variate Models
				Constant	$r(t)$	$s(t)$	
850	u	July	July	-11879.1	-11844.0	-11867.3	-11842.0
		Feb.	Feb.	-14370.0	-14367.8	-14316.7	-14315.3
		Feb.	July	-14597.0	-14553.6	-14429.2	-14492.6
		July	Feb.	-12046.1	-12022.4	-11942.7	-11954.1
850	v	July	July	-11925.2	-11873.3	-11875.4	-11853.2
		Feb.	Feb.	-14399.2	-14392.7	-14373.5	-14373.1
		Feb.	July	-14618.3	-14563.6	-14487.1	-14504.7
		July	Feb.	-12087.0	-12042.5	-11997.3	-11991.0
500	u	July	July	-13308.2	-13255.8	-13223.8	-13203.8
		Feb.	Feb.	-17944.6	-17889.7	-17890.6	-17853.6
		Feb.	July	-19419.3	-18592.2	-18411.8	-18170.3
		July	Feb.	-14143.8	-13857.6	-13788.7	-13633.3
500	v	July	July	-13314.7	-13274.6	-13274.8	-13253.6
		Feb.	Feb.	-17592.4	-17587.0	-17541.6	-17540.5
		Feb.	July	-18727.5	-18262.2	-17994.9	-17890.6
		July	Feb.	-13992.0	-13909.7	-13684.2	-13659.9
250	u	July	July	-17182.5	-17117.0	-17080.6	-17059.0
		Feb.	Feb.	-20872.6	-20836.3	-20538.2	-20505.4
		Feb.	July	-22345.1	-21676.2	-21016.2	-20887.3
		July	Feb.	-18057.7	-17829.7	-17266.9	-17178.3
250	v	July	July	-17005.0	-16900.6	-16889.1	-16833.3
		Feb.	Feb.	-20075.0	-20031.7	-19925.2	-19876.6
		Feb.	July	-20975.0	-20485.3	-20131.7	-19988.6
		July	Feb.	-17593.9	-17376.1	-17090.4	-16944.4

The values of $\tilde{\ell}$ for February data (respectively July data) using parameters fit using July data (respectively February data) are maximized most of the time by the two-variate model.

A comparison of the value of $\tilde{\ell}$, $\tilde{\ell}_c$, for the constant variance model of February (respectively July) data fit using the same month February (respectively July) data and the prediction values of $\tilde{\ell}$ for models (1)–(3) of February (respectively July) data using parameters estimated from the other month of July (respectively February) indicate the following. A majority of the time $\tilde{\ell}_c$ is larger than the corresponding values of $\tilde{\ell}$ for models (1)–(3) fit with the other month's data. This suggests that the first-guess covariate models fit using the other month's data may not describe the data as well as a constant variance model fit using the data being modeled. This may be an indication that models fit using first-guess February wind (respectively July wind) data are not good predictors of July (respectively February) wind component error.

Table 13 presents values of $\tilde{\ell}$ similar to those of Table 12 except that they are for the months of April and July. Comparison of the values of $\tilde{\ell}_c$ for data of one month fit with a constant variance model using the same data and the corresponding value of $\tilde{\ell}$ for the data using models with parameters estimated using the other month suggests that models using first-guess covariates do not have much predictive ability across these months. Table 14 presents the fraction of increase $(\tilde{\ell} - \tilde{\ell}_c) / |\tilde{\ell}_c|$ for the models with first guess covariates. Once again the results suggest that models using first guess wind components do not have much predictive ability across months.

**TABLE 13. VALUE OF LOG-LIKELIHOOD
FIRST-GUESS WIND COVARIATES
APRIL AND JULY**

Pressure Level	Wind Comp.	Data Set	Model	Constant	One-variate Models		Two-variate Models
					$r_f(t)$	$s_f(t)$	
850	u	July	July	-11879.1	-11844.0	-11867.3	<u>-11842.0</u>
		Apr.	Apr.	-14757.3	-14736.8	-14626.1	<u>-14625.9</u>
		Apr.	July	-14999.4	-14901.0	-14818.1	-14834.6
		July	Apr.	-12052.2	-11988.1	-11950.1	<u>-11945.1</u>
850	v	July	July	-11925.2	-11873.3	-11875.4	<u>-11853.2</u>
		Apr.	Apr.	-14949.1	-14937.9	-14848.1	<u>-14848.0</u>
		Apr.	July	-15247.5	-15169.4	-15002.4	-15022.3
		July	Apr.	-12133.8	-12077.1	-11992.8	-11996.6
500	u	July	July	-13308.2	-13255.8	-13223.8	<u>-13203.8</u>
		Apr.	Apr.	-17905.8	-17860.4	-17761.5	<u>-17742.2</u>
		Apr.	July	-18865.3	-18381.2	-18190.3	<u>-18031.5</u>
		July	Apr.	-13883.0	-13686.2	-13530.8	<u>-13499.0</u>
500	v	July	July	-13314.7	-13274.6	-13274.8	<u>-13253.6</u>
		Apr.	Apr.	-18112.7	-17948.5	-17703.9	<u>-17645.9</u>
		Apr.	July	-19233.9	-18557.9	-18350.3	<u>-18120.7</u>
		July	Apr.	-13967.8	-13597.8	-13465.5	<u>-13371.9</u>
250	u	July	July	-17182.5	-17117.0	-17080.6	<u>-17059.0</u>
		Apr.	Apr.	-22104.4	-22091.9	-22033.1	<u>-22001.7</u>
		Apr.	July	-23605.6	-23431.0	-22954.8	-22967.8
		July	Apr.	-18030.1	-18149.2	-17715.7	-17847.7
250	v	July	July	-17005.0	-16900.6	-16889.1	<u>-16833.3</u>
		Apr.	Apr.	-20637.7	-20576.6	-20355.8	<u>-20336.7</u>
		Apr.	July	-21139.5	-20837.5	-20503.3	<u>-20453.8</u>
		July	Apr.	-17346.8	-17149.6	-16965.5	<u>-16905.3</u>

TABLE 14. FRACTION OF INCREASE
IN LOG-LIKELIHOOD

$$(\bar{\ell} - \bar{\ell}_c) / |\bar{\ell}_c|$$

FIRST-GUESS WIND COVARIATES

Pressure Level	Wind Comp.	Data Set	Model	One-variate Models $r_f(t)$	One-variate Models $s_f(t)$	Two-variate Models
850	u	July	Feb.	*	*	*
		July	Apr.	*	*	*
		Feb.	July	*	*	*
		Apr.	July	*	*	*
850	v	July	Feb.	*	*	*
		July	Apr.	*	*	*
		Feb.	July	*	*	*
		Apr.	July	*	*	*
500	u	July	Feb.	*	*	*
		July	Apr.	*	*	*
		Feb.	July	*	*	*
		Apr.	July	*	*	*
500	v	July	Feb.	*	*	*
		July	Apr.	*	*	*
		Feb.	July	*	*	*
		Apr.	July	*	*	*
250	u	July	Feb.	*	*	0.00
		July	Apr.	*	*	*
		Feb.	July	*	*	*
		Apr.	July	*	*	*
250	v	July	Feb.	*	*	0.004
		July	Apr.	*	0.002	0.006
		Feb.	July	*	*	0.004
		Apr.	July	*	0.007	0.009

*: $\bar{\ell}_c$ (data described by model of constant variance fit using same data)

> $\bar{\ell}$ (data described by model fit using the other month)


4.3 Conclusions

Models (2) and (3) using observed wind components as covariates and fit using February or April (respectively July) data appear to have some predictive value for July (respectively February or April) data. Their predictive ability appears to be better for lower pressure levels. Models fit using April data appear to have more predictive ability than those fit using February data.

Models using first-guess wind covariates do not appear to have predictive ability across these months. It might be that models (1)–(3) fit with first-guess data from other Julys are better predictors of July wind component error. Alternatively, if first-guess winds are to be used as predictors, it might be worthwhile to develop a procedure to update the fitted model parameters using new data as it comes in.

850 MB U WIND; MODEL A ON DATA A; JULY OBS

1VAR=R[T]=0; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=0; 2VAR=+; BIN ON WS[T]





Figure 1


850 MB U WIND; MODEL B ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]




LOG AV PRED MSE PER BIN

Figure 2

850 MB U WIND; MODEL B ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 3

850 MB U WIND; MODEL A ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 4

850 MB V WIND; MODEL A ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 5

850 MB V WIND; MODEL B ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 6

850 MB V WIND; MODEL B ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 7

850 MB V WIND; MODEL A ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 8

500 MB U WIND; MODEL A ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 9

500 MB U WIND; MODEL B ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 10

500 MB U WIND; MODEL B ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 11

500 MB U WIND; MODEL A ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 12

500 MB V WIND; MODEL A ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 13

500 MB V WIND; MODEL B ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 14

500 MB V WIND; MODEL B ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 15

500 MB V WIND; MODEL A ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 16


250 MB U WIND; MODEL A ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]




LOG AV PRED MSE PER BIN

Figure 17

250 MB U WIND; MODEL B ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 18

250 MB U WIND; MODEL B ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 19

250 MB U WIND; MODEL A ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 20

250 MB V WIND; MODEL A ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 21

250 MB V WIND; MODEL B ON DATA B; JULY OBS

1VAR=R[T]=0; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=0; 2VAR=+; BIN ON WS[T]





Figure 22


250 MB V WIND; MODEL B ON DATA A; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]




LOG AV PRED MSE PER BIN

Figure 23


250 MB V WIND; MODEL A ON DATA B; JULY OBS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]



LOG AV PRED MSE PER BIN

Figure 24

REFERENCES

- M. Aitken (1987). "Modeling variance heterogeneity in normal regression using GLIM." *Appl. Stat.*, **36**, pp. 332-339.
- D. R. Cox and D. V. Hinkley, *Theoretical Statistics*, Chapman and Hall, London, 1974.
- M. Davidian and R. J. Carroll (1987) "Variance function estimation." *J. Am. Statis. Ass.*, **82**, pp. 1079-1091.
- J. S. Goerss and P. A. Phoebus (1991a). *The Multivariate Optimum Interpolation Analysis of Meteorological Data at FNOC*. NOARL Report Number 31, Naval Oceanographic and Atmospheric Research Laboratory, Stennis Space Center, MS.
- J. S. Goerss and P. A. Phoebus (1991b). "The Navy's operational atmospheric analysis." To appear in *Weather and Forecasting*.
- T. F. Hogan and T. E. Rosmond (1991) "The description of the Navy operational global atmospheric prediction system's spectral forecast model." *Monthly Weather Review*, **119**, No. 8, pp. 1786-1815.
- P. A. Jacobs and D. P. Gaver "Preliminary results from the analysis of wind component error," Naval Postgraduate School Technical Report NPSOR-91-029, September 1991.
- P. McCullagh and J. A. Nelder. *Generalized Linear Regression*, Chapman and Hall, New York, 1983.
- J. A. Nelder and Y. Lee. (1992). "Likelihood, quasi-likelihood and pseudolikelihood: some comparisons." *J. R. Statis. Soc. B.*, **54**, No. 1, pp. 273-284.

APPENDIX A

A BOOTSTRAP CROSS-VALIDATION STUDY FOR JULY DATA

In this Appendix histograms are presented from a bootstrap cross-validation study of models for July using both observed wind covariates and first guess wind covariates. Figures 1A-6A present results for the observed wind covariates. Figures 7A-12A present results for the first guess wind covariates.

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1 / 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA





Figure 1A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA





Figure 2A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA





Figure 3A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA





Figure 4A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1 / 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA





Figure 5A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA





Figure 6A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA; 1 G





Figure 7A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA;1 G





Figure 8A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA;1 G





Figure 9A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA; 1 G





Figure 10A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA;1 G





Figure 11A

FRAC INCR IN LIKE USING MODEL FIT WITH OTHER 1/ 2 DATA
 OVER CON VAR MOD FIT WITH SAME DATA;1 G





Figure 12A

APPENDIX B

A GRAPHICAL ASSESSMENT OF GOODNESS OF FIT AND CROSS- VALIDATION OF MODELS OF JULY WIND COMPONENT MEAN SQUARE ERROR USING FIRST-GUESS WIND COVARIATES

In this appendix we present figures assessing goodness of fit and cross-validation of the normal models (1)–(3) with first-guess wind covariates fit to July data. As in subsection (3.2) the data is randomly divided into two sets called DA and DB without regard to the values of the data; these sets are the same as those in that section.

The maximum likelihood parameter estimates for each model (1)–(3) are obtained for each set DA and DB and appear in Table 4. The estimated variances $\sigma_1^2(1,t)$, $\sigma_1^2(2,t)$, $\sigma_2^2(t)$ are computed for the parameters estimated from DA and DB using (1)–(3) for each data point in DA and DB.

To assess models (1) and (3) the data $(y(t), r(t), s(t))$ are binned into 10 bins based on ordering the values of $r(t)$ from smallest to largest. The data in the first bin correspond to the smaller values of $r(t)$; the data in the 10th bin correspond to the larger values of $r(t)$. Each bin contains about $\frac{1}{10}$ th of the data with the 10th bin containing a few more data. The averages of the estimated variances for models (1) and (3) are computed for each bin. The average $y(t)^2$ is also computed for each bin.

To assess models (2) and (3) the same procedure is used but the binning is based on values of $s(t)$.

Figures 1B-24B present graphs of the $\log[\text{average } y(t)^2]$ in each bin versus $\log[\text{average estimated variance}]$ in each bin for models (1) and (3) and models


(2) and (3). Figures 1B, 5B, 9B, 13B, 17B, 21B (respectively 2B, 6B, 10B, 14B, 18B, 22B) show the logarithm of the average of the $y(t)^2$ values of DA (respectively DB) versus the logarithm of the average of the estimated variances for each bin using the estimated parameters from DA (respectively DB). If a model were perfect, a point should be close to the 45° line shown. These figures assess goodness of fit.

Figures 3B, 7B, 11B, 15B, 19B, 23B (respectively 4B, 8B, 12B, 16B, 20B, 24B) present graphs of log average $y(t)^2$ of DA (respectively DB) versus log average estimated variances using parameters estimated using data DB (respectively DA). Once again if the model were perfect, the points would be close to the 45° line.

As suggested by the values of the log-likelihood $\tilde{\ell}$ in Tables 2 and 4, the figures for models using first-guess covariates indicate weaker goodness of fit and weaker cross-validation than Figures 1-24 for models with observed wind speed covariates. Both goodness-of-fit and cross-validation appear to improve somewhat for lower pressure levels; Figures 17B-24B. This suggests that models using first-guess covariates have somewhat better predictive and descriptive value at 250mb levels. However, they appear to be not as good as models using observed wind speed as covariates.

850 MB U WIND; MODEL A ON DATA A; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 1B

850 MB U WIND; MODEL B ON DATA B; JULY 1ST GUESS

1 VAR=R[T]=o; 2 VAR=+; BIN ON R[T]



1 VAR=WS[T]=o; 2 VAR=+; BIN ON WS[T]




Figure 2B

850 MB U WIND; MODEL B ON DATA A; JULY 1ST GUESS

1VAR=R[T]=°; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=°; 2VAR=+; BIN ON WS[T]





Figure 3B

850 MB U WIND; MODEL A ON DATA B; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 4B

850 MB V WIND; MODEL A ON DATA A; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 5B

850 MB V WIND; MODEL B ON DATA B; JULY 1ST GUESS

1VAR=R[T]=o;2VAR=+;BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o;2VAR=+;BIN ON WS[T]





Figure 6B

850 MB V WIND; MODEL B ON DATA A; JULY 1ST GUESS

1VAR=R[T]=o;2VAR=+;BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=o;2VAR=+;BIN ON WS[T]





Figure 7B

850 MB V WIND; MODEL A ON DATA B; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 8B

500 MB U WIND; MODEL A ON DATA A; JULY 1ST GUESS

1 VAR=R[T]=0;2 VAR=+;BIN ON R[T]



1 VAR=WS[T]=0;2 VAR=+;BIN ON WS[T]





Figure 9B

500 MB U WIND; MODEL B ON DATA B; JULY 1ST GUESS

1 VAR=R[T]=o;2 VAR=+;BIN ON R[T]



1 VAR=WS[T]=o;2 VAR=+;BIN ON WS[T]





Figure 10B

500 MB U WIND; MODEL B ON DATA A; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN
1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 11B

500 MB U WIND; MODEL A ON DATA B; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 12B

500 MB V WIND; MODEL A ON DATA A; JULY 1ST GUESS

1 VAR=R[T]=0;2 VAR=+;BIN ON R[T]



LOG AV PRED MSE PER BIN

1 VAR=WS[T]=0;2 VAR=+;BIN ON WS[T]





Figure 13B

500 MB V WIND; MODEL B ON DATA B; JULY 1ST GUESS

1 VAR=R[T]=:0;2 VAR=:+;BIN ON R[T]



1 VAR=WS[T]=:0;2 VAR=:+;BIN ON WS[T]





Figure 14B

500 MB V WIND; MODEL B ON DATA A; JULY 1ST GUESS

1 VAR=R[T]=o;2 VAR=+;BIN ON R[T]



1 VAR=WS[T]=o;2 VAR=+;BIN ON WS[T]





Figure 15B

500 MB V WIND; MODEL A ON DATA B; JULY 1ST GUESS

1 VAR=R[T]=0;2 VAR=+;BIN ON R[T]



1 VAR=WS[T]=0;2 VAR=+;BIN ON WS[T]





Figure 16B

250 MB U WIND; MODEL A ON DATA A; JULY 1ST GUESS

1 VAR=R[T]=0; 2 VAR=+; BIN ON R[T]



1 VAR=WS[T]=0; 2 VAR=+; BIN ON WS[T]





Figure 17B


250 MB U WIND; MODEL B ON DATA B; JULY 1ST GUESS

1 VAR=R[T]=0;2 VAR=+;BIN ON R[T]



LOG AV PRED MSE PER BIN

1 VAR=WS[T]=0;2 VAR=+;BIN ON WS[T]




LOG AV PRED MSE PER BIN

Figure 18B

250 MB U WIND; MODEL B ON DATA A; JULY 1ST GUESS

1 VAR=R[T]=o;2 VAR=+;BIN ON R[T]



1 VAR=WS[T]=o;2 VAR=+;BIN ON WS[T]





Figure 19B

250 MB U WIND; MODEL A ON DATA B; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



LOG AV PRED MSE PER BIN

1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]





Figure 20B

250 MB V WIND; MODEL A ON DATA A; JULY 1ST GUESS

1 VAR=R[T]=0;2 VAR=+;BIN ON R[T]



LOG AV PRED MSE PER BIN

1 VAR=WS[T]=0;2 VAR=+;BIN ON WS[T]





Figure 21B


250 MB V WIND; MODEL B ON DATA B; JULY 1ST GUESS

1 VAR=R[T]=0;2 VAR=+;BIN ON R[T]



LOG AV PRED MSE PER BIN

1 VAR=WS[T]=0;2 VAR=+;BIN ON WS[T]




LOG AV PRED MSE PER BIN

Figure 22B

250 MB V WIND; MODEL B ON DATA A; JULY 1ST GUESS

1 VAR=R[T]=0;2 VAR=+;BIN ON R[T]



1 VAR=WS[T]=0;2 VAR=+;BIN ON WS[T]





Figure 23B

250 MB V WIND; MODEL A ON DATA B; JULY 1ST GUESS

1VAR=R[T]=o; 2VAR=+; BIN ON R[T]



1VAR=WS[T]=o; 2VAR=+; BIN ON WS[T]




Figure 24B

INITIAL DISTRIBUTION LIST

1. Library (Code 52).....2
Naval Postgraduate School
Monterey, CA 93943-5000
2. Defense Technical Information Center.....2
Cameron Station
Alexandria, VA 22314
3. Office of Research Administration (Code 81)1
Naval Postgraduate School
Monterey, CA 93943-5000
4. Prof. Peter Purdue.....1
Code OR-Pd
Naval Postgraduate School
Monterey, CA 93943-5000
5. Department of Operations Research (Code 55).....1
Naval Postgraduate School
Monterey, CA 93943-5000
6. Prof. Donald Gaver, Code OR-Gv.....5
Naval Postgraduate School
Monterey, CA 93943-5000
7. Prof. Patricia Jacobs5
Code OR/Jc
Naval Postgraduate School
Monterey, CA 93943-5000
8. Center for Naval Analyses.....1
4401 Ford Avenue
Alexandria, VA 22302-0268
9. Dr. David Brillinger.....1
Statistics Department
University of California
Berkeley, CA 94720

10. Prof. Bernard Harris1
 Dept of Statistics
 University of Wisconsin
 610 Walnut Street
 Madison, WI 53706

11. Prof. W. R. Schucany.....1
 Dept. of Statistics
 Southern Methodist University
 Dallas, TX 75222

12. Prof. D. C. Siegmund.....1
 Dept. of Statistics
 Sequoia Hall
 Stanford University
 Stanford, CA 94305

13. Prof. H. Solomon1
 Department of Statistics
 Sequoia Hall
 Stanford University
 Stanford, CA 94305

14. Dr. Ed Wegman.....1
 George Mason University
 Fairfax, VA 22030

15. Dr. P. Welch1
 IBM Research Laboratory
 Yorktown Heights, NY 10598

16. Dr. Neil Gerr1
 Office of Naval Research
 Arlington, VA 22217

17. Prof. Roy Welsch.....1
 Sloan School
 M.I.T.
 Cambridge, MA 02139

18. Dr. J. Abrahams, Code 1111, Room 6071
 Mathematical Sciences Division, Office of Naval Research
 800 North Quincy Street
 Arlington, VA 22217-5000

19. Prof. J. R. Thompson.....1
 Dept. of Mathematical Science
 Rice University
 Houston, TX 77001

20. Prof. M. Leadbetter.....1
 Department of Statistics
 University of North Carolina
 Chapel Hill, NC 27514

21. Prof. J. B. Kadane.....1
 Dept. of Statistics
 Carnegie-Mellon University
 Pittsburgh, PA 15213

22. Prof. J. Lehoczky1
 Department of Statistics
 Carnegie-Mellon University
 Pittsburgh, PA 15213

23. Prof. M. Mazumdar1
 Dept. of Industrial Engineering
 University of Pittsburgh
 Pittsburgh, PA 15235

24. Prof. M. Rosenblatt.....1
 Department of Mathematics
 University of California, San Diego
 La Jolla, CA 92093

25. Prof. H. Chernoff.....1
 Department of Statistics
 Harvard University
 1 Oxford Street
 Cambridge, MA 02138

26. Prof. Frank Samaniego1
 Statistics Department
 University of California
 Davis, CA 95616

27.	Dr. James McKenna.....	1
	Bell Communications Research	
	445 South Street	
	Morristown, NJ 07960-1910	
28.	Prof. Brad Carlin.....	1
	School of Public Health	
	University of Minnesota	
	Mayo Bldg. A460	
	Minneapolis, MN 55455	
29.	Prof. W. Stuetzle.....	1
	Department of Statistics	
	University of Washington	
	Seattle, WA 98195	
30.	Prof. F. W. Mosteller.....	1
	Department of Statistics	
	Harvard University	
	1 Oxford St.	
	Cambridge, MA 02138	
31.	Dr. D. C. Hoaglin.....	1
	Department of Statistics	
	Harvard University	
	1 Oxford Street	
	Cambridge, MA 02138	
32.	Prof. N. D. Singpurwalla.....	1
	George Washington University	
	Washington, DC 20052	
33.	Prof. George S. Fishman.....	1
	Curr. in OR & Systems Analysis	
	University of North Carolina	
	Chapel Hill, NC 20742	
34.	Prof. Bradley Efron.....	1
	Statistics Dept.	
	Sequoia Hall	
	Stanford University	
	Stanford, CA 94305	

35. Prof. Carl N. Morris.....1
 Statistics Department
 Harvard University
 1 Oxford St.
 Cambridge, MA 02138
36. Dr. John E. Rolph.....1
 RAND Corporation
 1700 Main St.
 Santa Monica, CA 90406
37. Dr. Edward G. Coffman, Jr.....1
 AT&T Bell Telephone Laboratories
 600 Mountain Avenue
 Murray Hill, NJ 07974
38. Operations Research Center, Rm E40-164.....1
 Massachusetts Institute of Technology
 Attn: R. C. Larson and J. F. Shapiro
 Cambridge, MA 02139
39. Arthur P. Hurter, Jr.1
 Professor and Chairman
 Dept. of Industrial Engineering and Management Sciences
 Northwestern University
 Evanston, IL 60201-9990
40. Institute for Defense Analysis.....1
 1800 North Beauregard
 Alexandria, VA 22311
41. Prof. J. W. Tukey1
 Statistics Dept., Fine Hall
 Princeton University
 Princeton, NJ 08540
42. Dr. Colin Mallows.....1
 AT&T Bell Telephone Laboratories
 600 Mountain Avenue
 Murray Hill, NJ 07974

43. Dr. Jon Kettenring.....1
 Bellcore
 445 South Street
 Morris Township, NJ 07962-1910
44. Prof. David L. Wallace1
 Statistics Dept., University of Chicago
 5734 S. University Ave.
 Chicago, IL 60637
45. Dr. S. R. Dalal.....1
 Bellcore
 445 South Street
 Morristown, NJ 07962-1910
46. Dr. D. F. Daley1
 Statistic Dept. (I.A.S.)
 Australian National University
 Canberra, A.C.T. 2606
 AUSTRALIA
47. Dr. Guy Fayolle.....1
 I.N.R.I.A.
 Dom de Voluceau-Rocquencourt
 78150 Le Chesnay Cedex
 FRANCE
48. Professor H. G. Daellenbach.....1
 Department of Operations Research
 University of Canterbury
 Christchurch, NEW ZEALAND
49. Koh Peng Kong.....1
 OA Branch, DSO
 Ministry of Defense
 Blk 29 Middlesex Road
 SINGAPORE 1024
50. Professor Sir David Cox.....1
 Nuffield College
 Oxford, OX1 1NF
 ENGLAND

51. Dr. A. J. Lawrence.....1
 Dept. of Mathematics,
 University of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND
52. Dr. F. P. Kelly.....1
 Statistics Laboratory
 16 Mill Lane
 Cambridge
 ENGLAND
53. Dr. John Copas.....1
 Dept. of Mathematics,
 University of Birmingham
 P. O. Box 363
 Birmingham B15 2TT
 ENGLAND
54. Dr. D. Vere-Jones.....1
 Dept. of Math, Victoria Univ. of Wellington
 P. O. Box 196
 Wellington
 NEW ZEALAND
55. Prof. Guy Latouche1
 University Libre Bruxelles
 C.P. 212, Blvd. De Triomphe
 B-1050 Bruxelles
 BELGIUM
56. Dr. J. Goerss10
 Naval Oceanographic and Atmospheric Laboratory
 Monterey, CA 93943-5006

DUDLEY KNOX LIBRARY



3 2768 00330420 5